

- and prognosis in chronic systolic heart failure; the obesity paradox[J]. Am J Cardiol, 2003,91(7):891-894.
- [6] Oreopoulos A, Padwal R, Kalantar-Zadeh K, et al. Body mass index and mortality in heart failure: a meta-analysis[J]. Am Heart J, 2008,156(1): 13-22.
- [7] Fonarow GC, Srikanthan P, Costanzo MR, et al. An obesity paradox in acute heart failure: analysis of body mass index and in-hospital mortality for 108,927 patients in the acute decompensated heart failure national registry[J]. Am Heart J, 2007,153(1)3:74-81.
- [8] Lavie CJ, Milani RV, Ventura HO. Obesity and cardiovascular disease risk factor, paradox, and impact of weight loss[J]. J Am Coll Cardiol, 2009,53(21):1925-1932.
- [9] Horwich TB, Fonarow GC, Hamilton MA, et al. The relationship between obesity and mortality in patients with heart failure[J]. J Am Coll Cardiol, 2001,38(3):789-795.
- [10] Lavie CJ, Mehra MR, Milani RV. Obesity and heart failure prognosis: paradox or reverse epidemiology[J]. Eur Heart J, 2005, 26(1):5-7.
- [11] Mehra MR, Uber PA, Parh MH, et al. Obesity and suppressed B-type natriuretic peptide levels in heart failure[J]. J Am Coll Cardiol, 2004,43(9): 1590-1595.
- [12] Rauchhaus M, Coats AJS, Anker SD. The endotoxin-lipoprotein hypothesis[J]. Lancet, 2000,356(9233):930-933.
- [13] MacMahon SW, Wilcken DE, Macdonald GJ. The effect of weight reduction on left ventricular mass; a randomized controlled trial in young, overweight hypertensive patients[J]. N Engl J Med, 1986,314(6):334-339.
- [14] Alpert MA, Terry BE, Lambert CR, et al. Factors influencing left ventricular systolic function in nonhypertensive morbidly obese patients, and effect of weight loss induced by gastroplasty[J]. Am J Cardiol, 1993,71(8):733-737.
- [15] Sierra-Johnson J, Romero-Corral A, Somers VK, et al. Prognostic importance of weight loss in patients with coronary heart disease regardless of initial body mass index[J]. Eur Cardiovasc Prev Rehabil, 2008,15(3):336-340.
- [16] Pocock SJ, McMurray JJ, Dobson J, et al. Weight loss and mortality risk in patients with chronic heart failure in the candesartan in heart failure: assessment of reduction in mortality and morbidity(CHARM) programme[J]. Eur Heart J, 2008, 29(21): 2641-2650.
- [17] Mehra MR, Lavie CJ, Milani RV. Predicting prognosis in advanced heart failure: use of exercise indices[J]. Chest, 1996,110(2): 310-312.
- [18] Milani RV, Lavie CJ, Mehra MR, et al. Understanding the basics of cardiopulmonary exercise testing[J]. Mayo Clin Proc, 2006, 81(12):1603-1611.
- (收稿:2009-12-03)
(本文编辑:金谷英)

• 病例报告 •

隐匿性传导呈伪二度房室传导阻滞心电图分析 1 例

张 峰

DOI:10.3969/j.issn.1673-6583.2010.02.019

患者男性,50岁,有高血压病史,陈旧性下壁心肌梗死。心电图分析:P-R间期呈进行性延长,自0.26s开始逐次与QRS波脱离,虽然看似房室传导阻滞文氏现象,但是不见R-R间期进行性缩短,也不见有QRS波脱落,并伴有QRS波挫折,QRS时限0.11s。心率58次/分,绝大多数QRS与P波无关系,P-P间期、R-R间期相等。从梯形图看P-R间期自0.26s开始已经开始干扰了房室交界区的绝对不应期,使下传心室的P波受阻于此,并与窦性心搏相互干扰,故房室传导阻滞不成立。

讨论 心电图显示4相干扰性房室脱节,4相阻滞的产生与舒张电位自动除极化有较密切关系。由于心率减慢,舒张电位水平上移或阈电位水平上移,使窦性激动下传时受阻

于动作电位4相,以致在心电图中呈现隐匿性传导多形多样、单纯性或复杂的心律失常。



图1 患者心电图

作者单位:223900 江苏省宿迁市泗洪县人民医院功能科

(收稿:2009-02-18)

(本文编辑:丁媛媛)